

Unveiling the behavior of the ionized gas in irregular galaxies

Margarita Valdez-Gutiérrez

*Instituto Nacional de Astrofísica, Óptica y Electrónica, Tonantzintla,
 Puebla, México*

Margarita Rosado

Instituto de Astronomía-UNAM, México, City, México

1. Results and Discussion

We present the results of a detailed kinematical and dynamical analysis performed in two highly interesting gas-rich Irregular galaxies (Irrs): IC 1613 and NGC 4449. The analysis has been accomplished by means of optical Fabry–Perot interferometry mapping the $H\alpha$ and $[SII]$ lines. Our interest was centered on global and local scales in both galaxies. The results focused on the global radial velocity field in both galaxies show that they are different from each other. IC 1613 displays a highly perturbed velocity field due to local motions caused by superbubbles. NGC 4449 shows a velocity field which is relatively well behaved with a weak decreasing range in radial velocities ($\sim 60 \text{ km s}^{-1}$) from NE to SW. On local scales, we found that the HII region population for both galaxies displays a velocity dispersion mean value of $\sim 20 \text{ km s}^{-1}$, which does not reflect the environment of the host galaxy in each case. The HII region population in these galaxies display standard values according to their diameter distribution and luminosity functions ($D_0 = 30\text{--}65 \text{ pc}$; $\alpha = 1.5\text{--}1.9$). The superbubble population in IC 1613 displays expansion velocities ($20\text{--}40 \text{ km s}^{-1}$), sizes ($100\text{--}320 \text{ pc}$), dynamical ages ($0.7\text{--}2 \text{ Myr}$) and mechanical energies ($0.6\text{--}8 \times 10^{50} \text{ ergs}$) which can be explained –in the framework of the standard model– in terms of the activity derived from the massive stars (stellar winds and supernova explosions) at those locations. Based on our results (Valdez-Gutiérrez et al. 2001, 2002, Rosado et al. 2001), we conclude that the impressive distribution and kinematical/dynamical behavior of the ionized gas in these two gas-rich Irrs has been moulded –on global and local scales– by the activity of their massive stars; i.e. such activity has “etched” globally and locally the galaxies as they look today. Due to the nature of the present work –and as far as we know–, this is the first time that the kinematics and dynamics (on global and local scales) of the ionized gas in nearby gas-rich Irrs is unveiled in such detail.

References

- Valdez-Gutiérrez, M. et al. 2001, *A&A*, 366, 35
 Valdez-Gutiérrez, M. et al. 2002, *AJ*, submitted
 Rosado, M. et al. 2001, *AJ*, 122, 194